

HIGHER SECONDARY MODEL EXAMINATION 2012

HSE II

PART III

Max Score : 60

Time : 2 hrs

PHYSICS

General instruction to candidates

- Read all the questions carefully
- Total time for examination is 2 hrs 15 minutes including ‘cool off’ time
- First 15 minutes is cool off time during which neither answer the questions nor have discussion with others
- All questions are compulsory and only internal choice is allowed
- Choice is given for question numbered ‘6’

- 1) Two charges $+3 \mu\text{C}$ and $-3 \mu\text{C}$ are separated by a distance of 5 mm.
- a) What is the name of above configuration?
 - b) If the above arrangement is placed in a uniform electric field of intensity $3 \times 10^{-5} \text{N/C}$ with its axis perpendicular to the direction of the field, what is the torque acting on it?
 - c) If the arrangement is placed in a non-uniform electric field, what happens?
[1+2+1]
- 2) A cubical block is given a charge of $+40 \text{ mC}$ and placed in vacuum.
- a) Give the direction of electric field at the centre of each face.
 - b) Calculate the net flux through each face.
 - c) Will there be any change in flux if it is placed under water. Justify your answer.
(Dielectric constant of water = 81)
[1+2+1]
- 3) The emf of a dry cell is 1.5 V.
- a) What do you mean by emf?
 - b) Potentiometer is a better instrument than a voltmeter to measure emf of cell .why?
 - c) How will you use a potentiometer to compare the emf of two cells?
 - d) While measuring the emf, the primary circuit of the potentiometer should not change. Why?
[1+1+2+1]
- 4) A galvanometer is an instrument used for measuring small currents.
- a) What is the principle behind a galvanometer?
 - b) How is a galvanometer converted to ammeter and voltmeter?
 - c) Which has a higher resistance, millivoltmeter or microvoltmeter? Justify
[1+2+2]
- 5) Using Biot-savart’s law, we can find the magnetic field due to current distributions.
- a) State the above law
 - b) State another law which can be used for finding magnetic field due to current distributions
 - c) A torroid having inner and outer radius of 25 cm and 26 cm has 1000 turns carrying current of 5 A. What is the magnetic field inside and outside the core of torroid?
[1+1+2]
- 6) Faraday found that electric current can be generated with the help of magnetic field.
- a) State the laws of electromagnetic induction
 - b) Find an expression for self-inductance of a solenoid
 - c) Mention any two uses of eddy currents.
[2+2+1]
- OR**
- 6) A LCR series circuit can be used as tuning circuits.using resonance condition.
- a) Draw the corresponding phasor diagram.
 - b) Find the expression for resonant frequency.

- c) The current passing through inductor or capacitor only circuit is called wattless current. Why? [2+2+1]
- 7) A substance is weakly repelled by magnetic field
- Identify the substance. Give example
 - How is its susceptibility dependent on temperature?
 - Why is iron core preferred in transformer core [1+1+1]
- 8) A ray of light falls on one side of a prism whose refracting angle is A. The angles of incidence and refraction at the first face are i_1 and r_1 while at second face is i_2 and r_2 .
- Draw the incident and refracted ray at the minimum deviation.
 - What do you mean by critical angle of prism? If the refractive index of above prism is 1.5, find 'C'.
 - Find the angle of incidence in order that emergent ray may just graze the other side [1+2+1]
- 9) a) Which of the following phenomenon can't be explained by wave theory of light?
- Reflection
 - Refraction
 - Interference
 - Photoelectric effect
- b). In Young's experiment, the distance between slit and screen (D) = 1m, the separation of slits (d) = 1mm, wavelength of light (λ) = 600 nm. Calculate the fringe width (β)
- c) If the experiment is done in water ($n=4/3$) without changing the setup, what change will be observed in fringe width? Why? [1+2+2]
- 10) Electromagnetic spectrum consists of γ rays, X-rays, UV, Visible, IR, Microwaves and Radiowaves
- What is the phase difference between electric and magnetic field vectors in EM wave?
 - Which constituent radiation of the spectrum is mainly involved in
 - Greenhouse effect
 - Radar system
 - Distant photography during foggy condition
 - Treatment of cancer [1+2]
- 11) The resolving power of electron microscope is more than ordinary microscopes. Why?
- Which property of electrons is used in the construction of electron microscope?
 - Obtain expression of wavelength of De-Broglie waves associated with electrons accelerated through a potential of V volts. [1+2]
- 12) Classify the following statements into true or false and justify
- Radioactivity of a sample depends on the temperature of sample.
 - Rate of disintegration at any instant depends on the original number of atoms present initially [1+1]
- 13) The line spectrum of hydrogen was successfully explained by Bohr.
- What are the postulates of the above model?
 - Write the expression for energy and radius of orbit.
 - Find the longest wavelength in Lyman series and radius of second energy level. [1+2+2]
- 14) a) The lightly doped region of a transistor is known as
- b) Current amplification factor for CB, CE and CC transistor configuration are α , β and γ . Show that $\beta = \alpha / 1 - \alpha$.
- c) A transistor is having two PN Junctions. Is it possible to produce transistor action by fusing two PN junction diodes? Justify your answer [1+2+2]
- 15) a) An audio signal $10\sin 2\pi (1500t)$ amplitude modulates a carrier $40\sin 2\pi (100000t)$
- Sketch the AM Wave and what will be the percentage of modulation?
 - Calculate the area of region covered by TV broadcast by a tower of height 200m. (Radius of earth $R = 6.4 \times 10^6$ m) [2+2]